

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



3',5-Dihydroxy flavone 7-O- β -D-galacturonide-4'-O- β -D-glucopyranoside

FIG. 1




FIG. 2






FIG. 3



3',5-Dihydroxy flavone 7-O- β -D-galacturonide-4'-O- β -D-glucopyranoside

~~Fig 1~~
Fig. 1

Sample : 3',5-Dihydroxy flavone-7-O- β -D-galacturonide-
4'-O- β -D-glucopyranoside
Concentration : 0.0024 gm/10 mL H₂O
Inj. Vol. : 5 μ L
Column : RP-18, 5 μ m
Mobile Phase : 2% acetic acid in H₂O : ACN (83:17)
Flow rate : 1 mL/min.



~~Fig 2~~

Fig. 2

Sample : Fraction
Concentration : 0.8752 gm/10 mL H₂O
Inj. Vol. : 30 µ L
Column : RP-18, 5 µ m
Mobile Phase : 2% acetic acid in H₂O : ACN (83:17)
Flow rate : 1 mL/min.




Fig: 3

fig. 3